

REINWARDTIA

BEING A CONTINUATION OF THE

BULLETIN DU JARDIN BOTANIQUE DE BUITENZORG
(*BULLETIN OF THE BOTANIC GARDENS, BUITENZORG*)

EDITORS

A. DILMY

(Herbarium Bogoriense)

AND

C. G. G. J. VAN STEENIS

(Flora Malesiana)

Published by

HERBARIUM BOGORIENSE
KEBUN RAYA INDONESIA

ADDITIONAL NOTE ON CULLENIA CEYLANICA K. Schum. (Bombac).

A. J. G. H. KOSTERMANS *

Cullenia excelsa Wight in Pearson & Brown, *Commere. Timb. India* 1: 141, t. 51 1932; Foxworthy in *Philipp. J. Sci. Bot.* 4: 500. 1909; Trotter, *Common Comm. Timb. India* 65. 1901. Foxworthy in *Philip. J. Sci. Bot.* 4: 500. 1909.

Fruit collected in January 1958 from the tree cultivated in the Bogor Botanical Garden (no. XVI H. 13) enables me to correct and emendate the description published in "Communication of the Forest Research Institute, Bogor" no. 51 (1956) and in *Reinwardtia* 4 : 69.1956.

The ripe fruit are dirty green outside, they open on the branches and the seed falls out immediately.

Each fruit contains from one to 5 seeds, the common number, however, is one, the others do not develop; the fruit becomes pseudo-unilocular.

The chestnut brown seeds are completely enveloped by a lightbrown, somewhat glassy, slightly juicy and hardly sweet, furrowed aril of 2—4 mm thickness; the apical part of the aril is more or less fimbriate. — Figs. 38, 39.

*) D. Sc., Forest Research Institute, Bogor; Cooperator Herbarium Bogoriense, Bogor.




Fig. 38. — *Cullenia ceylanica* K. Schum.




Fig. 39. — *Cullenia ceylanica* K. Schum.; seeds with aril.